



Victor® Multi-Kill™ Place Pack

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ THIS LABEL and follow all use directions and use precautions.

IMPORTANT: Do not expose children, pets or other nontarget animals to rodenticides. To help prevent accidents:

1. Store product not in use in a location out of reach of children and pets.
2. Apply bait in locations out of reach of children, pets, domestic animals, and nontarget wildlife, or in tamper-resistant bait stations. These stations must be resistant to destruction by dogs and children under six years of age, and must be used in a manner that prevents such children from reaching into bait compartments and obtaining bait. If bait can be shaken from stations when they are lifted, units must be secured or otherwise immobilized. Even stronger bait stations are needed in areas open to hooved livestock, raccoons, bears, other potentially destructive animals, or in areas prone to vandalism.
3. Dispose of product container and any unused, spoiled or unconsumed bait as specified on this label.

USE RESTRICTIONS:

Do not use this product in sewers. Do not place bait near to or inside of ventilation duct openings. Do not contaminate water, food, feed, food or feed handling equipment, milk or meat handling equipment, or any area where there is a possibility of contaminating food, feed, milk, or meat, or surfaces that come into contact with food, feed, milk, or meat. All placements of this product in USDA-inspected facilities must be in tamper-resistant bait stations. Do not broadcast bait. Wear water-resistant gloves when handling bait.

Urban areas:

This product may be used in and around (within 15 feet of) homes, and in industrial, commercial, and public buildings. This product may also be used in transport vehicles (ships, trains, aircraft) and in and around (within 15 feet of) related port or terminal buildings. This product may be used in alleys.

Non-urban areas:

This product may only be used inside of homes and agricultural, industrial and commercial buildings.

SELECTION OF TREATMENT AREAS:

Determine areas where rats and mice will most likely find and consume the bait. Generally, these areas are along walls, by gnawed openings, beside burrows, in corners and concealed places, between floors and walls, or in locations where rats, mice, or their signs have been seen. Protect bait from rain and snow. Remove as much alternative food as possible.

APPLICATION DIRECTIONS:

Norway Rats and Roof Rats: Use 3 to 10 bait bags per placement, using tamper-resistant bait stations as required under "IMPORTANT". Space placements at intervals of 15 to 30 feet in infested areas pertaining to structures. Maintain an uninterrupted supply of fresh bait for at least 10 days or until fresh signs of feeding by rats cease to appear.

HOUSE MICE: Open bait bag and apply 1/4 to 1/2 ounce of bait per placement, using tamper-resistant bait stations as required under "IMPORTANT". Space placements at intervals of 8 to 12 feet in infested areas pertaining to structures. Larger quantities (up to one unopened 1.5 ounce bait bag) may be placed in areas of extremely high mouse activity. Maintain an uninterrupted supply of fresh bait for at least 15 days or until fresh signs of feeding by mice cease to appear.

FOLLOW-UP: Replace contaminated or spoiled bait immediately. Using water-resistant gloves, collect and properly dispose of dead animals and leftover bait, as well as any bait that has become accessible to non-target animals. To discourage re-infestation, limit sources of rodent food, water, and harborage as much as possible. If re-infestation does occur, repeat treatment. For continuous sources of infestation, set up permanent bait stations and replenish them as needed.

PRECAUTIONARY STATEMENTS:

Hazards To Humans and Domestic Animals

CAUTION: Harmful if swallowed. Wash thoroughly with soap and water after handling, and before eating, drinking, chewing gum, or using tobacco.

ENVIRONMENTAL HAZARDS:

This product is toxic to fish and wildlife. Predatory and scavenging mammals and birds might be poisoned if they feed upon animals that have eaten the bait. Do not apply directly to water.

STORAGE AND DISPOSAL:

Storage: Store in original container in a cool, dry place inaccessible to children and pets.

Disposal: If empty: Do not reuse this container. Place in trash or offer for recycling if available. If partly filled: Call your local solid waste agency instructions. Never place unused product down any indoor or outdoor drain.

FIRST AID:

Have label with you when obtaining treatment advice

If swallowed:

- Call a poison control center or doctor immediately for treatment advice.
- Have person sip a glass of water if able to swallow.
- Do not induce vomiting unless told to do so by the poison control center or doctor.

NOTE TO PHYSICIAN OR VETERINARIAN: If swallowed, this material may reduce the clotting ability of the blood and cause bleeding. If ingested, administer Vitamin K1 intramuscularly or orally. Repeat as necessary based on monitoring of prothrombin times.